

ANALYTICAL METHOD DEVELOPMENT FOR EXTRACTIVE SPECTROPHOTOMETRIC DETERMINATION OF Fe (II) AS A COMPLEX WITH BIS [3-HYDROXYIMINO - 5- METHYL - N-METHYL]-2-IMINE (HIMMI)

Rama S. Lokhande, Poonam P. Shevde and Sushama M. Lele

Department of Chemistry, University of Mumbai, Vidyanaagri, Santacruz (E),
Mumbai-400 098, India,

*E-mail: poonam_p_shevde@yahoo.co.in

ABSTRACT

Bis [3-Hydroxyimino - 5- Methyl - N- Methyl]-2-Imine (HIMMI) was synthesized and characterized by FTIR, NMR, elemental analysis as well as Mass spectrometry. This reagent was then applied for the development of the analytical method for the extractive spectrophotometric determination of Iron (II). Iron metal forms pinkish brown coloured complex, which can be extracted in chloroform at pH 6.0 having absorption maxima at 580 nm. Beer's law is obeyed in the concentration range 1-8 µg. The molar absorptivity and Sandell's Sensitivity of the extracted species are found to be $6.86 \times 10^3 \text{ Lit mol}^{-1} \text{ cm}^{-1}$ and $8.15 \times 10^{-3} \mu\text{g} / \text{cm}^2$ respectively. The developed method is highly sensitive, selective, simple, rapid, accurate, and has been satisfactorily applied for the determination of iron in the synthetic mixtures, pharmaceutical samples, and alloys.

Keywords: Characterization, Iron, Extractive Spectrophotometric determination.

© 2012 RASAYAN. All rights reserved.

INTRODUCTION

Iron is essential to nearly all known organisms. In cells, iron is generally stored in the centre of metalloproteins. The use of iron metal filings in organic synthesis is mainly for the reduction of nitro compounds.¹ Additionally, iron has been used for desulfurizations,² reduction of aldehydes,³ and the deoxygenation of amine oxides.⁴ Knowing the importance of Iron, a method highly sensitive, selective, simple, rapid, accurate and superior to reported methods in the literature⁵⁻³⁰ has been developed for its extractive spectrophotometric determination. The developed method has been satisfactorily applied for the determination of iron in the synthetic mixtures, pharmaceutical samples, and alloys.

EXPERIMENTAL

The HIMMI was synthesized³¹, characterized³¹ and used for extractive spectrophotometric determination of Fe(II). A stock solution of HIMMI (Concentration 0.005%) was prepared in methanol.

Iron (II) Solution

A weighed quantity of ferrous sulphate was dissolved in double distilled water containing dilute sulphuric acid and then diluted to the desired volume using double distilled water. The iron solution was then standardized by o-phenanthroline method³².

Recommended procedure

Mix 1 cm^3 aqueous solution containing 1-100 µg of iron and 1 cm^3 of 0.005 % methanolic solution of HIMMI in 25 cm^3 beaker. Adjust the pH of the solution to required value with dilute solution of H_2SO_4 and NaOH. Make the final aqueous volume up to 10 cm^3 . Transfer the solution into 125 cm^3 separating funnel and equilibrate for 1min with 10 cm^3 chloroform. Allow the two phases to separate and measure the absorbance of the organic extract containing the complex at 580 nm against reagent blank.

Table-1: Results and Discussion

Condition	Results
Absorption Maxima	580 nm
Solvent	Chloroform
pH range	5.5 – 6.5.
Equilibration time	1.0 min
Stability of Iron-HIMMI	12 h
Beer's range	1 to 8 $\mu\text{g}/\text{cm}^3$
Molar absorptivity	$6.86 \times 10^3 \text{ Lit mol}^{-1} \text{ cm}^{-1}$
Sandell's sensitivity	$8.15 \times 10^{-3} \mu\text{g}/\text{cm}^2$
Mole Ratio of Fe : HIMMI	1 :2




Fig.-1: Slope Ratio Method




Fig.-2: Calibration Plot

Effect of foreign ions

The effect of diverse ions on the iron (II) determination was studied, in presence of a definite amount of a foreign ion. Various cations and anions were investigated in order to find the tolerance limit of these foreign ions in the extraction of iron (II) (Table 2). The tolerance limit of the foreign ion was taken as the amount required causing an error of not more than +2% in the recovery of Iron (II). The ions which interfere in the spectrophotometric determination of iron were masked by using appropriate masking agents (Table 3).

Table-2: Effect of foreign ions

S. No.	Interfering ions	Tolerance Limit (mg)
1	Acetate, Oxalate, Citrate, SCN^- , $\text{S}_2\text{O}_7^{2-}$, NO_3^- , Urea, NO_2^- , $\text{S}_2\text{O}_8^{2-}$, SO_4^{2-} , SO_3^{2-} , IO_3^- , BrO_3^- , ClO_3^- , Cl^- , Br^- , I^- , F^- .	10
2	Phosphate,	8
3	Thiourea	10
4	Zr(II), Ba(II), Mo(VI), Mg(II), Zn(II),	10
5	Ca(II), Al(III), Ce(IV), Mn(II)	5
6	Tl(I), Ag(I), Th(IV), Cr(VI), Li(I),	5
7	U(VI)	4
8	V(VI), Cd(II), Pb(II),	2
9	Rh(III), As(III)	1
10	Co^{+2} , Cu^{+2} , Ni^{+2} , Pt^{+2} , Pd^{+2} , EDTA, Hg^{+2}	Interfere strongly

Applications

The present method was applied for determination of amount of Iron (II) in various samples as alloys, synthetic mixtures, injection vial, tablets and industrial waste water. The results obtained were well in agreement with those of standard methods, Table-4.

Table-3: Effect of masking agent

Interfering Ion	Masking agent added	Interfering Ion	Masking agent added
Ag(I)	Potassium iodide	Citrate	Sodium molybdate
Cd(II)	Potassium iodide	Tartarate	Sodium molybdate
Pb(II)	Sodium thiosulphate	EDTA	Boiling with concentrated HNO ₃
Mn(II)	Sodium fluoride	CN ⁻	Boiling with concentrated HNO ₃ and formaldehyde
Ce(IV)	Sodium fluoride	Zr(IV)	Sodium fluoride
Cr(II)	Ammonium acetate	Ni(II)	Thiourea

Table-4: Applications

S. No.	Sample	Amount of Fe(II)	
		Standard Method	Present Method
1.	Alloy /Ore		
(a)	Hematite	35.0%	34.98%
(b)	Steel	67.2%	66.99%
2.	Capsule/tablet		
(a)	Injection Viol	50 mg/ml	49.9 mg/ml
(b)	Vit B12 (Capsule)	1.5 mg	1.49 mg
3.	Synthetic Mixture		
(a)	Fe(II)(5)+Zn(II)(5)	4.98 ppm	4.99 ppm
(b)	Fe(II)(5)+Mg(II)(5)	4.98 ppm	4.99 ppm
4.	Industrial Waste Water at Ulhasnagar Creek	0.5 ppm	0.49 ppm

Every result is average of three independent determinations.

CONCLUSION

The results obtained show that the newly developed method in which the reagent **HIMMI** was used, can be effectively used for quantitative extraction and estimation Fe (II) from aqueous media. The proposed method is quick and requires less volume of organic solvent. The results show good agreement with the standard methods. The method is very precise, faster and simpler than other methods. The method is precise, accurate, less time consuming and easily employed anywhere, even in small laboratories as it requires only uv visible spectrophotometer and not much sophisticated and costly measurement devices or instrumentation.

REFERENCES

1. B. A. Fox and T. L. Threlfall, *Organic Syntheses.*, **5**, 346 (1973).
2. A. T. Blomquist and L. I. Dinguid, *J. Org. Chem.*, **12**, 718, (1947).
3. H. T. Clarke and E. E. Dreger, *Org. Syn.*, **1**,304, (1941).
4. J.Den Hertog and Overhoff, *J. Recl. Trav. Chim.* **69**, 468, (1950).
5. C. H. R. Gentry and L. S. Sherrington, *Analyst.*, **75**, 17, (1950).
6. D. J.Eve and A. J. S. Strasheim, *Afr. Chem. Inst.*, **9**, 5, (1956).
7. Y. Tanaka and Ito. K. Japan, *Analyst.*, **6**, 728, (1957).
8. M. Oosting, *Anal. Khim acta.*, **21**, 397, (1959).
9. T. Sekine and D. J. Dryssen, *Inorg. Nuclear Chem.*, **26**, 2013, (1964).
10. H. A. Mttola and H. Freiser, *Talanta.*, **13**,55(1966).
11. J. R. Jhonston and W. J. Halland, *Mikrochim Acta.*, **1**, 126, (1972).

12. M. Valcarcel, D. Perz Bendito and F.Pin., *Perez.AnQuim.*, **25**,1, (1972).
13. M. Valcarcel and F. Pin, *Perez.AnQuim.*,**68**, 383, (1972).
14. R.B. Singh, B.S. garg and R.P. Singh,*Talanta.*, **26**, 425, (1975).
15. C. K. Bhaskar and S. Devi., *Talanta.*,**25**, 5449, (1979).
16. T. S. Reddy and S.B. Rao., *Curr.Sci.*, **48**, 439, (1979).
17. D. Yamakato, S. Hirqoka and M. Hikawa. *Bunseki kagaku.*, **30**, 626, (1981).
18. V. R.Patil, R. B. Kharatand, B. K. Deshmukh., *J. Inorg. Nucl. Chem.*, **43**, 3397, (1981).
19. D. Novova, and K. Stoyanov., *Anal.Chim.Acta.*, **138**, 321, (1982).
20. A. K. Chakrabarti., *Ind.J.Chem.*, **21**, 439, (1982).
21. T. Nakanishi and M. Otomo., *Nippon Kagaku Kaishi.*, **4**, 518, (1983).
22. K. Lal and S. R. Malhotra, *J. Ind. Chem. Soc.*, **60**, 308, (1983).
23. S. Q. R. Iyes and A. P. Joshi., *Ind. J. Chem.*, **221**, 907, (1983).
24. A. K. Malik and A. L. Rao, *Ind. J. Chem.*, **329 A**, 829, (1983).
25. H. S. Gowda and S. M. Ahmed., *Ind. J. Chem.*, **22**, 1086, (1983).
26. A. K. Chakrabarti., *Ind.J. Chem.*, **25**, 886, (1986).
27. S. K. Kuchekar, M. A. Aruze and M. B. Chavan., *Ind.J. Chem.*, **25**, 1041, (1986).
28. P. Sarkar, P. K. Karia and S. K. Muzumdar, *Ind. J. Chem. Soc.*, **26**, 987, (1987).
29. S. P. Arya., *Talanta.*, **34**, 293, (1987).
30. D. M. Rao, K. H. Reddy, D. V. Reddy. And Shrikrishnadevaraya. *Ind.J Chem. SecA*, 1145, (1989).
31. R. S. Lokhande, V. R. Patil, P. P. Shevde and S. M. Lele ., *Int. J. Chem. Sci.*, **8(1)**, 88, (2010)
32. A. I. Vogel; 'Textbook of Quantitative chemical Analysis', 5th Ed.E.L.B.S.(1991).

[RJC-907/2012]

International Journal of
Chemical, Environmental and Pharmaceutical Research

www.ijcepr.com

ISSN: 2229-3892(Print); ISSN: 2229-5283(Online)

[Abstracted in : Chemical Abstracts Service , American Chemical Society, USA and CAB(I) , UK]

ijCEPr widely covers all fields of **Chemical, Environmental and Pharmaceutical Research.**

Manuscript Categories: Full-length paper, Review Articles, Short/Rapid Communications.

Manuscripts should be addressed to:

E-mail: ijcepr@gmail.com

Water: Research & Development [Water R&D]

www.waterrnd.com

ISSN: 2249-2003

[Abstracted in : Chemical Abstracts Service, USA and CAB(I) , UK]

WaterR&D is an international Research Journal, dedicated to 'Water'. It is a truly interdisciplinary journal on water science and technology. It'll showcase the latest research related to Water in the field of chemistry, physics, biology, agricultural, food, pharmaceutical science, and environmental, oceanographic, and atmospheric science. It includes publication of reviews, regular research papers, case studies, communications and short notes.

Manuscript Categories: Full-length paper, Review Articles, Short/Rapid Communications.

Manuscripts should be addressed to:

E-mail: waterrnd@gmail.com